

Underdiagnosis and undertreatment of modifiable cardiovascular risk factors: a Childhood Cancer Survivor Study (CCSS) report

Background: Survivors of childhood cancer exposed to cardiotoxic therapies are at increased risk of heart disease. Hypertension, dyslipidemia, and diabetes are modifiable cardiovascular risk factors (CVRFs) that synergistically increase this risk. Therefore, we aimed to determine the prevalence of and predictors associated with CVRF underdiagnosis and undertreatment in this population.

Methods: CCSS participants at increased risk of heart disease due to prior cancer therapy were enrolled in an ongoing randomized intervention trial (NCT03104543) to improve CVRF identification and treatment. Participants completed a baseline survey (CVRF status, lifestyle habits, attitudes towards healthcare), anthropometry, and blood draw. Blood pressure, low density lipoprotein, triglyceride, glucose and Hgb A1c were measured and classified as normal/abnormal per standard clinical criteria. Multivariable logistic regression estimated odds ratios (OR [95% confidence intervals]) associated with predictors and risk of CVRF underdiagnosis and undertreatment.

Results: As of January 2020, 522 participants (43% male) were available for analysis (47% response), with a median age 38y (range 20-65) and 28y (18-49) from original cancer treatment (75% anthracycline, 47% chest radiation). With mean measured BMI 27.3 ± 6.5 kg/m², self-reported prevalence rates were hypertension 27%, dyslipidemia 33%, and diabetes 9%. While 90% of participants had a routine check-up ≤ 2 y ago, 58% had a measured CVRF in the abnormal range. Specifically, among previously undiagnosed participants, we observed rates of abnormal blood pressure (26%), lipids (17%), and glucose tolerance (27%). Among those with pre-existing hypertension, dyslipidemia, and diabetes, 11%, 49%, and 54%, respectively, had measurements outside of the usual therapeutic target range. In multivariable analysis, BMI ≥ 25 kg/m² (vs < 25) was associated with risk of underdiagnosis (OR 1.8 [1.2-2.8]). For undertreatment, significant adverse factors included older age (> 35 vs ≤ 35 y: OR 2.5 [1.2-5.1]), BMI ≥ 30 kg/m² (vs < 25 : OR 3.3 [1.7-6.4]), and greater perceived reliance on others for healthcare decisions (OR 1.7 [1.2-2.4]). Those with greater health-related self-efficacy were less likely to be undertreated (OR 0.5 [0.3-0.96]).

Conclusions: CVRF underdiagnosis and undertreatment among childhood cancer survivors at increased risk of heart disease was common. Greater awareness among survivors and primary care providers and more aggressive control of CVRFs may mitigate this risk.

Eric J. Chow^{1,2,3}
Yan Chen⁴
Gregory T. Armstrong⁵
Laura-Mae Baldwin⁶
Todd M. Gibson⁷
Melissa M. Hudson^{5,8}
Aaron McDonald⁵
Paul C. Nathan⁹
Karen Syrjala^{1,2}
Emily S. Tonorezos¹⁰
Kevin C. Oeffinger¹¹
Yutaka Yasui^{4,7}

¹Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle WA

²Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle WA

³Department of Pediatrics, Seattle Children's Hospital, University of Washington, Seattle, WA

⁴University of Alberta, Edmonton, Alberta, Canada

⁵Department of Epidemiology and Cancer Control, St. Jude Children's Research Hospital, Memphis, TN

⁶Department of Family Medicine, University of Washington, Seattle, WA

⁷National Cancer Institute, Bethesda, MD

⁸Department of Oncology, St. Jude Children's Research Hospital, Memphis, TN

⁹Department of Pediatrics, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada

¹⁰Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY

¹¹Department of Medicine, Duke University, Durham, NC